

The availability of Blue-fronted Amazon nesting cavities is at risk in Central-West Brazil

Gláucia H. F. Seixas* and Guilherme Mourão
Corresponding author: glauucia@fundacaoneotropica.org.br

Appendix 1. Number of dead and alive trees with suitable cavities for nesting Blue-fronted Amazon found in the study area from 2016 to 2024 and its percentage in relation to the grand total.

Species	Local name	Dead	Alive	Total	%
<i>Acrocomia aculeata</i>	Macaúba	1724	0	1724	95.1
<i>Albizia niopoides</i>	Farinha seca	0	2	2	0.1
<i>Anadenanthera colubrina</i>	Angico branco	0	3	3	0.2
<i>Aspidosperma sp</i>	Peroba	0	1	1	0.1
<i>Caesalpinia peltophoroides</i>	Sibipiruna	0	3	3	0.2
<i>Campomanesia xanthocarpa</i>	Guabiroba	0	7	7	0.4
<i>Copaifera Langsdorffii</i>	Copaiba	0	2	2	0.1
<i>Gallesia intergrifolia</i>	Pau d'alho	0	2	2	0.1
<i>Grevillea robusta</i>	Grevílea	0	1	1	0.1
<i>Handroanthus avellaneda</i>	Ipê roxo	1	2	3	0.2
<i>Mauritia flexuosa</i>	Buriti	2	0	2	0.1
<i>Peltophorum dubium</i>	Canafistula	0	5	5	0.3
<i>Persea americana</i>	Abacate	0	2	2	0.1
<i>Sapindus saponaria</i>	Saboneteira	0	1	1	0.1
<i>Syagrus romanzoffiana</i>	Pindó	27	0	27	1.5
Not identified	NI	27	0	27	1.5
Grand total		1781	31	1812	100